Reference Determination Content

Example document — fictitious patient. A suggested draft determination prepared by the Veterans Health Centre from the matching example Diagnostic Assessment, to show how a DVA delegate could determine the claim. Names, dates, service details and clinicians are invented. It is not a decision of the Department of Veterans' Affairs.

Left Ankle - Osteoarthritis

Evidence

The evidence that I have considered in making this decision includes:

- Formal Claim with supporting statements;
- A Statement of your Service from the Department of Defence;
- Copies of your Service Medical Records.
- A Diagnostic Assessment by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 3 November 2025;
- A Diagnosis Form (D9287) by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 3 November 2025;
- An entry medical examination by the army recruiting medical officer, dated 18 August 1949;
- A repatriation medical examination of returned prisoner of war by the medical officer, dated 6 September 1953;
- An X-ray left ankle (emergency record and operation report) by the radiologist and orthopaedic surgeon, dated 14 March 1962;
- An operation report (left knee meniscectomy) by the orthopaedic surgeon, dated 8 May 1963;
- A field hospital admission and operation report by the surgeon, Vietnam, dated 21 September 1966;
- An outpatient health record and pathology by the medical officer, dated 9 July 1969;
- An endocrinology report (absorptiometry and long-bone survey) by the endocrinologist and radiologist, dated 17 November 1972;
- An X-ray left tibia and pathology by the radiologist, dated 14 February 1973;
- X-ray pelvis by the radiologist, dated 11 March 1974;
- A periodic health examination by the medical officer, dated 20 October 1975;
- An outpatient health record and X-ray left ankle by the medical officer and radiologist, dated 2 June 1976;
- A specialist report by the orthopaedic surgeon, dated 10 August 1976;
- A dual-photon absorptiometry report by the radiologist, dated 22 October 1984;

- A specialist report and X-ray left ankle by the orthopaedic surgeon and radiologist, dated 18 June 1985;
- An operation report (left ankle arthrodesis) by the orthopaedic surgeon, dated 3 March 1986.

Your service

I have identified that you have the following service:
Australian Defence Force service between :

18 August 1949 and 11 April 1987

Left Ankle - Osteoarthritis

You have stated that your condition was caused by:

- *having inflammatory joint disease as specified, of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical onset of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical onset of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical onset of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical onset of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having increased bone mineral density before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at least two years before the clinical onset of osteoarthritis in that joint; and (b) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical onset of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical onset of osteoarthritis in that joint*

- *having acromegaly before the clinical onset of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical onset of osteoarthritis in that joint*

Diagnosis: Osteoarthritis

Basis of diagnosis: The diagnosis was made in the Diagnostic Assessment by Dr Thomas Perkins dated 3 November 2025, and is supported by a specialist report by the orthopaedic surgeon, dated 10 August 1976.

I am satisfied this is the appropriate diagnosis.
In determining your claim for this condition, I have applied the SOP:

Statement of Principles concerning Osteoarthritis (Balance of Probabilities) (No. 62 of 2017)

which sets out the factors that can connect the condition with service.

I am satisfied that you have met the following factor:

- *having inflammatory joint disease as specified, of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical onset of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical onset of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical onset of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical onset of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having increased bone mineral density before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at least two years before the clinical onset of osteoarthritis in that joint; and (b) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical onset of osteoarthritis in that joint*

- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having acromegaly before the clinical onset of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical onset of osteoarthritis in that joint*

I have noted your contention as to the cause of this condition. The available evidence indicates, I have noted your contention to the cause of your condition. Your service medical records indicate that you presented to the health facility with mechanical pain in the left ankle on weight bearing, worse on ladders, stairs and uneven ground, morning stiffness of about 20 minutes, swelling after activity, loss of ankle movement and a worsening limp on 2 June 1976.

The diagnosis was confirmed by weight-bearing X-ray of the left ankle on 2 June 1976, which showed joint space narrowing, marginal osteophytes and subchondral sclerosis of the tibiotalar joint, and by an orthopaedic surgeon on 10 August 1976. This meets the Statement of Principles definition of osteoarthritis, a degenerative joint disease characterised by loss of articular cartilage and new bone formation.

This has then led to further investigations and the diagnosis of your condition of *Osteoarthritis*. I have accepted your condition of *Osteoarthritis*.

Having considered your service records and the available medical evidence, I am satisfied the SOP is met and that this condition is related to your peacetime service.

I have therefore accepted that this condition is related to your service with effect from 15 May 1976.

Left Ankle - Osteoarthritis (Worsening)

Evidence

The evidence that I have considered in making this decision includes:

- Formal Claim with supporting statements;
- A Statement of your Service from the Department of Defence;
- Copies of your Service Medical Records.
- A Diagnostic Assessment by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 3 November 2025;
- A Diagnosis Form (D9287) by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 3 November 2025;
- An entry medical examination by the army recruiting medical officer, dated 18 August 1949;
- A repatriation medical examination of returned prisoner of war by the medical officer, dated 6 September 1953;
- An X-ray left ankle (emergency record and operation report) by the radiologist and orthopaedic surgeon, dated 14 March 1962;
- An operation report (left knee meniscectomy) by the orthopaedic surgeon, dated 8 May 1963;
- A field hospital admission and operation report by the surgeon, Vietnam, dated 21 September 1966;
- An outpatient health record and pathology by the medical officer, dated 9 July 1969;
- An endocrinology report (absorptiometry and long-bone survey) by the endocrinologist and radiologist, dated 17 November 1972;
- An X-ray left tibia and pathology by the radiologist, dated 14 February 1973;
- X-ray pelvis by the radiologist, dated 11 March 1974;
- A periodic health examination by the medical officer, dated 20 October 1975;
- An outpatient health record and X-ray left ankle by the medical officer and radiologist, dated 2 June 1976;
- A specialist report by the orthopaedic surgeon, dated 10 August 1976;
- A dual-photon absorptiometry report by the radiologist, dated 22 October 1984;
- A specialist report and X-ray left ankle by the orthopaedic surgeon and radiologist, dated 18 June 1985;
- An operation report (left ankle arthrodesis) by the orthopaedic surgeon, dated 3 March 1986.

Your service

I have identified that you have the following service:
Australian Defence Force service between :

18 August 1949 and 11 April 1987

Left Ankle - Osteoarthritis (Worsening)

You have stated that the worsening of your condition was caused by:

- *having inflammatory joint disease as specified, of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical worsening of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical worsening of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical worsening of*

osteoarthritis in that joint

- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at two years before the clinical worsening of osteoarthritis in that joint; and (b) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical worsening of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having acromegaly before the clinical worsening of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical worsening of osteoarthritis in that joint*

- *inability to obtain appropriate clinical management for osteoarthritis*

Diagnosis: Osteoarthritis (Worsening)

Basis of diagnosis: The diagnosis was made in the Diagnostic Assessment by Dr Thomas Perkins dated 3 November 2025, and is supported by a specialist report by the orthopaedic surgeon, dated 10 August 1976.

I am satisfied this is the appropriate diagnosis.
In determining your claim for this condition, I have applied the SOP:

Statement of Principles concerning Osteoarthritis (Balance of Probabilities) (No. 62 of 2017)

which sets out the factors that can connect the condition with service.

I am satisfied that you have met the following factor:

- *having inflammatory joint disease as specified, of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical worsening of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical worsening of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical worsening of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at two years before the clinical worsening of osteoarthritis in that joint; and (b) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical worsening of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical worsening of osteoarthritis in that joint*

- *having Paget's disease of bone of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having acromegaly before the clinical worsening of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical worsening of osteoarthritis in that joint*
- *inability to obtain appropriate clinical management for osteoarthritis*

I have noted your contention as to the cause of this condition. The available evidence indicates, I have noted your contention to the cause of your condition. Your service medical records indicate that you presented to the health facility with mechanical pain in the left ankle on weight bearing, worse on ladders, stairs and uneven ground, morning stiffness of about 20 minutes, swelling after activity, loss of ankle movement and a worsening limp on 2 June 1976.

The diagnosis was confirmed by weight-bearing X-ray of the left ankle on 2 June 1976, which showed joint space narrowing, marginal osteophytes and subchondral sclerosis of the tibiotalar joint, and by an orthopaedic surgeon on 10 August 1976. This meets the Statement of Principles definition of osteoarthritis, a degenerative joint disease characterised by loss of articular cartilage and new bone formation.

After the condition arose, it was worsened during your service. The reactive arthritis of the left ankle of 1952 and 1960 preceded the documented worsening on 18 June 1985. The Staphylococcus aureus septic arthritis of the left ankle of September 1966 preceded the worsening on 18 June 1985. The intra-articular trimalleolar fracture of 14 March 1962 preceded the worsening on 18 June 1985. The haemarthrosis of 1962 preceded the worsening, and a further haemarthrosis was aspirated after a fall from a ladder on 4 August 1983, during service. Gout of the left ankle continued after onset, with attacks in 1978, 1981 and 1984, before the worsening on 18 June 1985. The 1962 fracture and the fall from a ladder on 4 August 1983, which caused a haemarthrosis and was treated by immobilisation, both occurred within the 25 years before the worsening, from 18 June 1960 to 18 June 1985. The acute osteochondral tear of the talar dome of 14 March 1962 occurred 23 years before the worsening, within the 25-year window. The acute medial meniscal tear of the left knee of 8 May 1963 occurred within the 25 years before the worsening. The frostbite of the left foot and ankle of 1951 to 1952 preceded the worsening. The varus and rotational malunion persisted and progressed to a fixed varus deformity, present throughout the five years before the worsening, from 18 June 1980 to 18 June 1985. The dysbaric osteonecrosis of the left talar dome, shown in 1972, preceded the worsening and was associated with collapse of the talar dome on the 1985 X-ray. Your heavy gripping work as an Army diver for about 35 hours a week from 1958 to 1974 preceded the worsening. You used pneumatic percussive tools on more days than not from 1960 to 1974, more than ten years within the 25 years before the worsening. The right toe amputations and asymmetric gait, present since 1952, continued throughout the five years before the worsening. The lifting of about 1 500 000 kg from 1958 to 1968 preceded the worsening, which occurred in 1985, within the 25 years

following that period. The carrying of about 4 800 hours from 1958 to 1968 preceded the worsening, which occurred within the 25 years following that period. The ladder climbing of about 300 rungs a day from 1960 to 1974, a continuous period of 14 years within the 25 years before the worsening, preceded it. Your BMI remained above 25 from 1962 to the worsening in 1985, and was 30.2 in June 1985. Your waist exceeded 102 cm from 1964 to the worsening, and was 109 cm in June 1985. Your waist exceeded 102 cm, and therefore also 88 cm, for more than ten years before the worsening. The kneeling and squatting of about two hours a day from 1958 to 1974 preceded the worsening, which occurred within the 25 years following that period. The internal derangement of the left knee from 1963 persisted before the worsening. The left chondromalacia patella diagnosed in 1964 persisted before the worsening. The beriberi neuropathy with loss of proprioception at the left ankle persisted before the worsening. Paget's disease of the distal left tibia remained active, with a raised alkaline phosphatase in 1984, before the worsening. Acromegaly, treated surgically in 1972, preceded the worsening. Lateral instability of the left ankle persisted from 1963 to the worsening, including the giving way that caused the fall of 4 August 1983. The left hip femoroacetabular impingement shown in 1974 persisted before the worsening. An orthopaedic surgeon recommended a custom brace in 1976 and arthrodesis of the left ankle in 1981, but the brace was not supplied for two years and the arthrodesis was deferred for four years by postings and the military hospital waiting list, while you continued to climb ladders as an instructor.

This has then led to further investigations and the diagnosis of your condition of *Osteoarthritis (Worsening)*. I have accepted your condition of *Osteoarthritis (Worsening)*.

Having considered your service records and the available medical evidence, I am satisfied the SOP is met and that the worsening of this condition is related to your peacetime service.

I have therefore accepted that the worsening of this condition is related to your service with effect from 18 June 1985.

Example document — fictitious patient. A suggested draft determination prepared by the Veterans Health Centre from the matching example Diagnostic Assessment, to show how a DVA delegate could determine the claim. Names, dates, service details and clinicians are invented. It is not a decision of the Department of Veterans' Affairs.