Reference Determination Content

Example document — fictitious patient. A suggested draft determination prepared by the Veterans Health Centre from the matching example Diagnostic Assessment, to show how a DVA delegate could determine the claim. Names, dates, service details and clinicians are invented. It is not a decision of the Department of Veterans' Affairs.

Left Knee - Medial Femoral Condyle - Osteoarthritis

Evidence

The evidence that I have considered in making this decision includes:

- Formal Claim with supporting statements;
- A Statement of your Service from the Department of Defence;
- Copies of your Service Medical Records.
- A Diagnostic Assessment by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 6 November 2025;
- A Diagnosis Form (D9287) by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 6 November 2025;
- An inpatient record and X-ray report by the naval hospital, dated 3 March 1988;
- A diving medicine record by the medical officer, Persian Gulf, dated 16 March 1991;
- An operation report by the surgeon, deployed hospital, East Timor, dated 12 December 1999;
- An inpatient record (deployed hospital) by the iraq, dated 31 March 2003;
- An MRI left knee by the radiologist, dated 15 April 2003;
- An operation report by the orthopaedic surgeon, dated 28 April 2003;
- A rheumatology consultation by the rheumatologist, dated 6 May 2004;
- An MRI left knee and femur by the radiologist, dated 7 June 2005;
- A haematology and endocrinology review by the haematologist and endocrinologist, dated 22 September 2006;
- An outpatient health record by the medical officer, Afghanistan, dated 15 January 2009;
- An X-ray left knee by the radiologist, dated 12 May 2009;
- An outpatient health record by the medical officer, naval base health centre, dated 21 July 2009;
- An X-ray left knee by the radiologist, dated 24 July 2009;
- A bone densitometry report by the nuclear medicine physician, dated 21 November 2011;
- An outpatient health record and aspiration note (role 2 facility) by the afghanistan, dated 18 June 2012;
- An X-ray and MRI left knee by the radiologist, dated 11 October 2012;

- An operation report by the orthopaedic surgeon, dated 14 May 2013.

Your service

I have identified that you have the following service:
Australian Defence Force service between :

14 January 1980 and 11 December 2014

Left Knee - Medial Femoral Condyle - Osteoarthritis

You have stated that your condition was caused by:

- *having inflammatory joint disease as specified, of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical onset of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical onset of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical onset of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical onset of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having increased bone mineral density before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at least two years before the clinical onset of osteoarthritis in that joint; and (b) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical onset of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having acromegaly before the clinical onset of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical onset of osteoarthritis in that joint*

- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical onset of osteoarthritis in that joint*

Diagnosis: Medial Femoral Condyle - Osteoarthritis

Basis of diagnosis: The diagnosis was made in the Diagnostic Assessment by Dr Thomas Perkins dated 6 November 2025, and is supported by an X-ray left knee by the radiologist, dated 24 July 2009.

I am satisfied this is the appropriate diagnosis.
In determining your claim for this condition, I have applied the SOP:

Statement of Principles concerning Osteoarthritis (Balance of Probabilities) (No. 62 of 2017)

which sets out the factors that can connect the condition with service.

I am satisfied that you have met the following factor:

- *having inflammatory joint disease as specified, of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical onset of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical onset of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical onset of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an*

average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical onset of osteoarthritis in that joint

- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having increased bone mineral density before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at least two years before the clinical onset of osteoarthritis in that joint; and (b) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical onset of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical onset of osteoarthritis in that joint*

- *having Paget's disease of bone of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having acromegaly before the clinical onset of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical onset of osteoarthritis in that joint*

I have noted your contention as to the cause of this condition. The available evidence indicates, I have noted your contention to the cause of your condition. Your service medical records indicate that you presented to the health facility with aching pain on the inner side of the left knee on walking, climbing ladders and standing, morning stiffness of about 20 minutes, grating and intermittent swelling on 21 July 2009.

The diagnosis was confirmed by weight-bearing X-rays of the left knee on 24 July 2009, which showed narrowing of the medial tibiofemoral joint space with marginal osteophytes and subchondral sclerosis of the medial femoral condyle, Kellgren-Lawrence grade 2, in varus alignment. This meets the Statement of Principles definition of osteoarthritis.

This has then led to further investigations and the diagnosis of your condition of *Medial Femoral Condyle - Osteoarthritis*. I have accepted your condition of *Medial Femoral Condyle - Osteoarthritis*.

Having considered your service records and the available medical evidence, I am satisfied the SOP is met and that this condition is related to your peacetime service.

I have therefore accepted that this condition is related to your service with effect from 14 July 2009.

Left Knee - Medial Femoral Condyle - Osteoarthritis (Worsening)

Evidence

The evidence that I have considered in making this decision includes:

- Formal Claim with supporting statements;
- A Statement of your Service from the Department of Defence;
- Copies of your Service Medical Records.
- A Diagnostic Assessment by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 6 November 2025;
- A Diagnosis Form (D9287) by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 6 November 2025;
- An inpatient record and X-ray report by the naval hospital, dated 3 March 1988;
- A diving medicine record by the medical officer, Persian Gulf, dated 16 March 1991;
- An operation report by the surgeon, deployed hospital, East Timor, dated 12 December 1999;
- An inpatient record (deployed hospital) by the iraq, dated 31 March 2003;
- An MRI left knee by the radiologist, dated 15 April 2003;
- An operation report by the orthopaedic surgeon, dated 28 April 2003;
- A rheumatology consultation by the rheumatologist, dated 6 May 2004;
- An MRI left knee and femur by the radiologist, dated 7 June 2005;
- A haematology and endocrinology review by the haematologist and endocrinologist, dated 22 September 2006;
- An outpatient health record by the medical officer, Afghanistan, dated 15 January 2009;
- An X-ray left knee by the radiologist, dated 12 May 2009;
- An outpatient health record by the medical officer, naval base health centre, dated 21 July 2009;
- An X-ray left knee by the radiologist, dated 24 July 2009;
- A bone densitometry report by the nuclear medicine physician, dated 21 November 2011;
- An outpatient health record and aspiration note (role 2 facility) by the afghanistan, dated 18 June 2012;
- An X-ray and MRI left knee by the radiologist, dated 11 October 2012;
- An operation report by the orthopaedic surgeon, dated 14 May 2013.

Your service

I have identified that you have the following service:
Australian Defence Force service between :

14 January 1980 and 11 December 2014

Left Knee - Medial Femoral Condyle - Osteoarthritis (Worsening)

You have stated that the worsening of your condition was caused by:

- *having inflammatory joint disease as specified, of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical worsening of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical worsening of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical worsening of osteoarthritis in that*

joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period

- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at two years before the clinical worsening of osteoarthritis in that joint; and (b) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical worsening of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having acromegaly before the clinical worsening of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical worsening of osteoarthritis in that joint*
- *inability to obtain appropriate clinical management for osteoarthritis*

Diagnosis: Medial Femoral Condyle - Osteoarthritis (Worsening)

Basis of diagnosis: The diagnosis was made in the Diagnostic Assessment by Dr Thomas Perkins dated 6 November 2025, and is supported by an X-ray left knee by the radiologist, dated 24 July 2009.

I am satisfied this is the appropriate diagnosis.
In determining your claim for this condition, I have applied the SOP:

Statement of Principles concerning Osteoarthritis (Balance of Probabilities) (No. 62 of 2017)

which sets out the factors that can connect the condition with service.

I am satisfied that you have met the following factor:

- *having inflammatory joint disease as specified, of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical worsening of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical worsening of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical worsening of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at two years before the clinical worsening of osteoarthritis in that joint; and (b) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical worsening of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having acromegaly before the clinical worsening of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical worsening of osteoarthritis in that joint*

- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical worsening of osteoarthritis in that joint*
- *inability to obtain appropriate clinical management for osteoarthritis*

I have noted your contention as to the cause of this condition. The available evidence indicates, I have noted your contention to the cause of your condition. Your service medical records indicate that you presented to the health facility with aching pain on the inner side of the left knee on walking, climbing ladders and standing, morning stiffness of about 20 minutes, grating and intermittent swelling on 21 July 2009.

The diagnosis was confirmed by weight-bearing X-rays of the left knee on 24 July 2009, which showed narrowing of the medial tibiofemoral joint space with marginal osteophytes and subchondral sclerosis of the medial femoral condyle, Kellgren-Lawrence grade 2, in varus alignment. This meets the Statement of Principles definition of osteoarthritis.

After the condition arose, it was worsened during your service. Psoriatic arthropathy flared with a left knee effusion in August 2011, before the worsening on 11 October 2012. The bacterial septic arthritis of the left knee in December 1999 preceded the worsening. The intra-articular osteochondral fracture of the medial femoral condyle from March 2003 preceded the worsening. On 18 June 2012 a twisting fall in Afghanistan caused a haemarthrosis of the left knee, with 60 millilitres of blood aspirated, before the worsening. Haemochromatosis arthropathy and chondrocalcinosis of the left knee persisted, with an acute pseudogout attack in 2011, before the worsening. The 25 years before the worsening run from 11 October 1987 to 11 October 2012. MRI on 11 October 2012 showed an acute chondral flap tear of the medial femoral condyle from the fall of 18 June 2012, within the 25 years before the worsening. The fall of 18 June 2012 caused an acute radial tear of the remnant medial meniscus, shown on MRI on 11 October 2012, within the 25 years before the worsening. The frostbite of the left knee in January 2009 preceded the worsening. The varus malunion of the left femur persisted for more than twenty years before the worsening. The dysbaric osteonecrosis of the medial femoral condyle progressed to subchondral collapse, shown on MRI on 11 October 2012, before the worsening. The forceful gripping work from 1982 to 1997 amounted to more than ten years within a continuous 15-year period before the worsening. Your use of vibrating percussive tools on most working days from 1987 to 1998, eleven years, fell within the 25 years before the worsening, which run from 11 October 1987 to 11 October 2012. The asymmetric gait from the femoral shortening persisted for more than twenty years before the worsening. The lifting of more than 200,000 kilograms from 1985 to 1995 was followed by the worsening in 2012, within the 25 years following that period. The carrying of more than 4,500 hours from 1984 to 1994 was followed by the worsening in 2012, within the 25 years following that period. The ladder and stair climbing from 1987 to 1996, nine continuous years, fell within the 25 years before the worsening. Your BMI remained between 28 and 31 throughout the ten years before the worsening. Your waist circumference exceeded 102 centimetres throughout the ten years before the worsening, measured at 109 centimetres in 2011. Your waist circumference exceeded 88 centimetres throughout the ten years before the worsening. The ten years of kneeling from 1984 to 1994 were followed by the

worsening in 2012, within the 25 years following that period. Internal derangement of the left knee, including the meniscal tears of 2003 and 2012, preceded the worsening. Chondromalacia patellae of the left knee, diagnosed in 1992, preceded the worsening. The loss of proprioception in both legs from spinal decompression sickness persisted before the worsening. Paget's disease of the left distal femur remained active before the worsening. Acromegaly preceded the worsening. Instability of the left knee from the 2003 cruciate tear persisted, and the knee gave way causing the fall of 18 June 2012, more than one year before the worsening. Pincer-type femoroacetabular impingement of the left hip preceded the worsening. After the injury of 18 June 2012 in Afghanistan you had a swollen, locking knee, but no MRI, physiotherapy or orthopaedic review was available at the forward base and your request for aeromedical evacuation for assessment was not approved.

This has then led to further investigations and the diagnosis of your condition of *Medial Femoral Condyle - Osteoarthritis (Worsening)*. I have accepted your condition of *Medial Femoral Condyle - Osteoarthritis (Worsening)*.

Having considered your service records and the available medical evidence, I am satisfied the SOP is met and that the worsening of this condition is related to your peacetime service.

I have therefore accepted that the worsening of this condition is related to your service with effect from 11 October 2012.

Example document — fictitious patient. A suggested draft determination prepared by the Veterans Health Centre from the matching example Diagnostic Assessment, to show how a DVA delegate could determine the claim. Names, dates, service details and clinicians are invented. It is not a decision of the Department of Veterans' Affairs.