Reference Determination Content

Example document — fictitious patient. A suggested draft determination prepared by the Veterans Health Centre from the matching example Diagnostic Assessment, to show how a DVA delegate could determine the claim. Names, dates, service details and clinicians are invented. It is not a decision of the Department of Veterans' Affairs.

Right Knee - Osteoarthritis

Evidence

The evidence that I have considered in making this decision includes:

- Formal Claim with supporting statements;
- A Statement of your Service from the Department of Defence;
- Copies of your Service Medical Records.
- A Diagnostic Assessment by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 18 March 2025;
- A Diagnosis Form (D9287) by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 18 March 2025;
- A casualty record and X-ray report by the hospital ship, Persian Gulf, dated 3 March 1991;
- An operation report by the orthopaedic surgeon, dated 6 March 1991;
- An MRI right knee by the radiologist, dated 20 June 1994;
- An arthroscopy operation report by the orthopaedic surgeon, dated 2 August 1994;
- An inpatient record by the infectious diseases physician, dated 11 August 1994;
- A joint aspiration and pathology report by the medical officer, Naval base health centre, dated 11 March 1997;
- A rheumatology consultation by the rheumatologist, dated 22 July 1998;
- A field hospital record by the surgeon, East Timor, dated 21 November 1999;
- MRI long bones by the radiologist, dated 15 November 2002;
- A recompression treatment record by the diving medical officer, dated 8 April 2003;
- A neurology consultation by the neurologist, dated 16 June 2003;
- A periodic health examination and diving medical by the medical officer, dated 22 August 2005;
- A field medical record by the medical officer, Afghanistan, dated 19 January 2006;
- An outpatient health record by the medical officer, Naval base health centre, dated 25 October 2006;
- An X-ray right knee by the radiologist, dated 1 November 2006;
- A bone densitometry report by the endocrinologist, dated 16 August 2010;
- An X-ray and MRI right knee by the radiologist, dated 3 October 2011;

- An orthopaedic consultation by the orthopaedic surgeon, dated 12 October 2011.

Your service

I have identified that you have the following service:
Australian Defence Force service between :

20 January 1990 and 15 June 2021

Right Knee - Osteoarthritis

You have stated that your condition was caused by:

- *having inflammatory joint disease as specified, of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical onset of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical onset of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical onset of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical onset of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having increased bone mineral density before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at least two years before the clinical onset of osteoarthritis in that joint; and (b) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical onset of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having acromegaly before the clinical onset of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical onset of osteoarthritis in that joint*

- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical onset of osteoarthritis in that joint*

Diagnosis: Osteoarthritis

Basis of diagnosis: The diagnosis was made in the Diagnostic Assessment by Dr Thomas Perkins dated 18 March 2025, and is supported by an X-ray right knee by the radiologist, dated 1 November 2006.

I am satisfied this is the appropriate diagnosis.
In determining your claim for this condition, I have applied the SOP:

Statement of Principles concerning Osteoarthritis (Balance of Probabilities) (No. 62 of 2017)

which sets out the factors that can connect the condition with service.

I am satisfied that you have met the following factor:

- *having inflammatory joint disease as specified, of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical onset of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical onset of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical onset of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical onset of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical onset of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an*

average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical onset of osteoarthritis in that joint

- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical onset of osteoarthritis in that joint; and (ii) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having increased bone mineral density before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at least two years before the clinical onset of osteoarthritis in that joint; and (b) where the clinical onset of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical onset of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical onset of osteoarthritis in that joint*

- *having Paget's disease of bone of the affected joint before the clinical onset of osteoarthritis in that joint*
- *having acromegaly before the clinical onset of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical onset of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical onset of osteoarthritis in that joint*

I have noted your contention as to the cause of this condition. The available evidence indicates, I have noted your contention to the cause of your condition. Your service medical records indicate that you presented to the health facility with aching right knee pain, morning stiffness of about 30 minutes, pain on ladders and kneeling, and swelling after diving days on 25 October 2006.

The diagnosis was confirmed by X-rays of the right knee on 1 November 2006, which showed new medial and lateral joint space narrowing, marginal osteophytes and subchondral sclerosis, Kellgren-Lawrence grade 2, where X-rays of 22 August 2005 had shown preserved joint spaces. With pain, stiffness and impaired function, this meets the Statement of Principles definition of osteoarthritis: a degenerative joint disorder with clinical manifestations of pain, impaired function and stiffness and radiological evidence of loss of articular cartilage and osteophytes.

This has then led to further investigations and the diagnosis of your condition of *Osteoarthritis*. I have accepted your condition of *Osteoarthritis*.

Having considered your service records and the available medical evidence, I am satisfied the SOP is met and that this condition is related to your peacetime service.

I have therefore accepted that this condition is related to your service with effect from 18 October 2006.

Right Knee - Osteoarthritis (Worsening)

Evidence

The evidence that I have considered in making this decision includes:

- Formal Claim with supporting statements;
- A Statement of your Service from the Department of Defence;
- Copies of your Service Medical Records.
- A Diagnostic Assessment by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 18 March 2025;
- A Diagnosis Form (D9287) by Dr Thomas Perkins, General Practitioner, Veterans Health Centre, dated 18 March 2025;
- A casualty record and X-ray report by the hospital ship, Persian Gulf, dated 3 March 1991;
- An operation report by the orthopaedic surgeon, dated 6 March 1991;
- An MRI right knee by the radiologist, dated 20 June 1994;
- An arthroscopy operation report by the orthopaedic surgeon, dated 2 August 1994;
- An inpatient record by the infectious diseases physician, dated 11 August 1994;
- A joint aspiration and pathology report by the medical officer, Naval base health centre, dated 11 March 1997;
- A rheumatology consultation by the rheumatologist, dated 22 July 1998;
- A field hospital record by the surgeon, East Timor, dated 21 November 1999;
- MRI long bones by the radiologist, dated 15 November 2002;
- A recompression treatment record by the diving medical officer, dated 8 April 2003;
- A neurology consultation by the neurologist, dated 16 June 2003;
- A periodic health examination and diving medical by the medical officer, dated 22 August 2005;
- A field medical record by the medical officer, Afghanistan, dated 19 January 2006;
- An outpatient health record by the medical officer, Naval base health centre, dated 25 October 2006;
- An X-ray right knee by the radiologist, dated 1 November 2006;
- A bone densitometry report by the endocrinologist, dated 16 August 2010;
- An X-ray and MRI right knee by the radiologist, dated 3 October 2011;
- An orthopaedic consultation by the orthopaedic surgeon, dated 12 October 2011.

Your service

I have identified that you have the following service:
Australian Defence Force service between :

20 January 1990 and 15 June 2021

Right Knee - Osteoarthritis (Worsening)

You have stated that the worsening of your condition was caused by:

- *having inflammatory joint disease as specified, of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical worsening of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical worsening of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*

- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at two years before the clinical worsening of osteoarthritis in that joint; and (b) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical worsening of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having acromegaly before the clinical worsening of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical worsening of osteoarthritis in that joint*
- *inability to obtain appropriate clinical management for osteoarthritis*

Diagnosis: Osteoarthritis (Worsening)

Basis of diagnosis: The diagnosis was made in the Diagnostic Assessment by Dr Thomas Perkins dated 18 March 2025, and is supported by an X-ray right knee by the radiologist, dated 1 November 2006.

I am satisfied this is the appropriate diagnosis.
In determining your claim for this condition, I have applied the SOP:

Statement of Principles concerning Osteoarthritis (Balance of Probabilities) (No. 62 of 2017)

which sets out the factors that can connect the condition with service.

I am satisfied that you have met the following factor:

- *having inflammatory joint disease as specified, of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having an infection of the affected joint as specified before the clinical worsening of osteoarthritis in that joint*
- *having an intra-articular fracture of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having haemarthrosis of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having a depositional joint disease as specified, in the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having trauma to the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute articular cartilage tear of the affected joint within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having an acute meniscal tear of the affected knee within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *having frostbite involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having disordered joint mechanics of the affected joint for at least five years before the clinical worsening of osteoarthritis in that joint*
- *having necrosis of the subchondral bone near the affected joint, including that from dysbaric osteonecrosis, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the upper limb only, performing any combination of repetitive activities or tasks involving repeated or sustained pinch grip or hand/power grip, for an average of at least 30 hours per week, for a cumulative period of at least ten years within a continuous period of 15 years, before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the hand, wrist or elbow joint only, using a hand-held, vibrating percussive tool or object on more days than not, for a cumulative period of at least ten years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb only: having: (i) an amputation involving either leg; or (ii) an asymmetric gait; for at least five years before the clinical worsening of osteoarthritis in that joint*

- *for osteoarthritis of a joint of the lower limb only: lifting loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 150 000 kilograms within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: carrying loads of at least 20 kilograms while bearing weight through the affected joint: (i) to a cumulative total of at least 3 800 hours within any ten year period before the clinical worsening of osteoarthritis in that joint; and (ii) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a joint of the lower limb only: ascending or descending at least 150 stairs or rungs of a ladder per day, on more days than not, for a continuous period of at least five years, within the 25 years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: being overweight for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for males, having a waist circumference exceeding 102 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a joint of the lower limb or hand joint only: for females, having a waist circumference exceeding 88 centimetres for at least ten years before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip or knee joint only: (a) kneeling or squatting for a cumulative period of at least one hour per day, on more days than not, for a continuous period of at two years before the clinical worsening of osteoarthritis in that joint; and (b) where the clinical worsening of osteoarthritis in that joint occurs within the 25 years following that period*
- *for osteoarthritis of a knee joint only, having internal derangement of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of the patello-femoral joint only, having chondromalacia patella before the clinical worsening of osteoarthritis in that joint*
- *having a disorder associated with loss of pain sensation or proprioception involving the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having Paget's disease of bone of the affected joint before the clinical worsening of osteoarthritis in that joint*
- *having acromegaly before the clinical worsening of osteoarthritis in that joint*
- *having joint instability or dislocation of the affected joint, at least one year before the clinical worsening of osteoarthritis in that joint*
- *for osteoarthritis of a hip joint only, having femoroacetabular impingement syndrome of the affected joint, before the clinical worsening of osteoarthritis in that joint*
- *inability to obtain appropriate clinical management for osteoarthritis*

I have noted your contention as to the cause of this condition. The available evidence indicates, I have noted your contention to the cause of your condition. Your service medical records indicate that you presented to the health facility with aching right knee pain, morning stiffness of about 30 minutes, pain on ladders and kneeling, and swelling after diving days on 25 October 2006.

The diagnosis was confirmed by X-rays of the right knee on 1 November 2006, which showed new medial and lateral joint space narrowing, marginal osteophytes and subchondral sclerosis, Kellgren-Lawrence grade 2, where X-rays of 22 August 2005 had shown preserved joint spaces. With pain, stiffness and impaired function, this meets the Statement of Principles definition of osteoarthritis: a degenerative joint disorder with clinical manifestations of pain, impaired function and stiffness and radiological evidence of loss of articular cartilage and osteophytes.

After the condition arose, it was worsened during your service. The psoriatic arthropathy of the right knee remained active, with flares in 2008 and 2010 treated with prednisolone, and was present before the worsening documented on 3 October 2011. The septic arthritis of August 1994 affected the right knee and preceded the worsening in 2011. The intra-articular osteochondral fracture of June 1994 preceded the worsening, and MRI on 3 October 2011 showed collapse of the previously fractured medial femoral condyle surface. On 14 June 2011 the ship's medical officer aspirated 40 mL of blood from the right knee after the fall on a ship's ladder. You had a further attack of gout of the right knee in March 2010, with urate crystals on aspiration, before the worsening. The fall of 14 June 2011 applied significant force through the right knee, with immediate pain, swelling and loss of movement lasting more than seven days. MRI on 3 October 2011 showed a new chondral flap tear of the medial femoral condyle from the fall of 14 June 2011, and the 1994 chondral tear also lay within the 25 years before the worsening. MRI on 3 October 2011 showed a fresh medial meniscal tear from the fall of 14 June 2011, within the 25 years before the worsening. The frostbite involving the right knee in January 2006 preceded the worsening in 2011. The 7 degrees of valgus from the 1991 malunion persisted throughout the five years before the worsening, from 3 October 2006 to 3 October 2011, and increased to 10 degrees on standing films in October 2011. The dysbaric osteonecrosis of the distal right femur was still present on MRI on 3 October 2011 and preceded the worsening. Your gripping duties continued at 32 to 40 hours a week from 1996 to 2011, a cumulative 15 years within the continuous 15-year period before the worsening. Daily use of impact wrenches, needle guns and chipping hammers continued from 1992 to 2011, a cumulative 19 years within the 25 years before the worsening. The asymmetric antalgic gait persisted and worsened after onset, and was present throughout the five years before the worsening, from 3 October 2006 to 3 October 2011. Between October 2001 and September 2011 you continued to lift diving sets, charges and boarding equipment of 20 to 40 kg while standing, more than 1 500 000 kilograms within that ten-year period, and the worsening fell within the 25 years following it. Between October 2001 and September 2011 you carried diving sets and boarding equipment of 20 kg or more for more than 3 800 hours, and the worsening fell within the 25 years following that period. Ladder climbing of more than 150 rungs a day on more days than not continued from 1993 to 2011, including the five years before the worsening, all within the 25 years before it. Your BMI remained between 28 and 30 from 2001 to 2011, so you were overweight throughout the ten

years before the worsening, from 3 October 2001 to 3 October 2011. You are male, and your waist circumference remained above 102 cm, recorded at 108 cm in 2011, throughout the ten years before the worsening. You are male, and your waist circumference exceeded 88 centimetres throughout the ten years before the worsening. Kneeling for at least one hour a day on more days than not continued from 1990 to 2011, more than two years, and the worsening fell within the 25 years following that period. The graft laxity and the fresh meniscal tear of June 2011 are internal derangement of the right knee present before the worsening. Chondromalacia of the right patella persisted and preceded the worsening; The sensory and proprioceptive loss over the right leg from the 2003 decompression illness persisted, confirmed on neurological review in 2009, before the worsening. Paget's disease of the right proximal tibia persisted, with a raised alkaline phosphatase in 2010, before the worsening. Acromegaly, treated in 2003, preceded the worsening. Recurrent giving way of the right knee continued from 1996 and caused the fall of 14 June 2011. The right femoroacetabular impingement syndrome diagnosed in 2004 persisted before the worsening. After the fall on warlike service in the Middle East on 14 June 2011 you could not obtain imaging, orthopaedic review or physiotherapy at sea and continued boarding duties in a brace for 12 weeks.

This has then led to further investigations and the diagnosis of your condition of *Osteoarthritis (Worsening)*. I have accepted your condition of *Osteoarthritis (Worsening)*.

Having considered your service records and the available medical evidence, I am satisfied the SOP is met and that the worsening of this condition is related to your peacetime service.

I have therefore accepted that the worsening of this condition is related to your service with effect from 3 October 2011.

Example document — fictitious patient. A suggested draft determination prepared by the Veterans Health Centre from the matching example Diagnostic Assessment, to show how a DVA delegate could determine the claim. Names, dates, service details and clinicians are invented. It is not a decision of the Department of Veterans' Affairs.